



MP Biomedicals, LLC

29525 Fountain Parkway  
Solon, Ohio 44139

Telephone: 440/337-1200  
Toll Free: 800/854-0530  
Fax: 440/337-1180  
mailto: [biotech@mpbio.com](mailto:biotech@mpbio.com)  
web: <http://www.mpbio.com>

## TECHNICAL INFORMATION

cappel

**PEROXIDASE-CONJUGATED GOAT IGG FRACTION TO  
HUMAN SECRETORY IGA - 2ML**  
Catalog #: 55215  
Lot #: 07532

Total Protein: 5.3 mg/ml  
Titer: 1:194,000

### DESCRIPTION:

Product is the lyophilized powder of horseradish peroxidase (HRP)-conjugated goat IgG fraction to human secretory IgA and buffer salts.

Antibody and highly purified HRP ( $R_z > 3.0$ ) are conjugated under defined conditions to obtain optimally labeled product. Conjugated protein is purified by salt fractionation. The product is dialyzed into 0.02M sodium phosphate, 0.14M sodium chloride, pH 7.3, adjusted to standard titer, filtered through a 0.22  $\mu$ m filter, vialled and lyophilized. No preservative has been added.

### USES:

This product is suitable for use as a reagent in enzyme immunoassays (EIA), cell and tissue staining (for light microscopy), cell and tissue labeling (for electron microscopy), and blot immunostaining. (Note:  $F(ab')_2$  fragments are recommended for staining of cells or tissues which contain Fc receptors. Affinity purified antibodies are recommended to avoid non-specific binding from inherent antibodies of host animals. See **Related Product Ordering Information** for these products.)

### SPECIFICATIONS:

The total protein is measured using the Biuret procedure with bovine albumin as the standard. To ensure that the product titer falls within the required range, antibody titer is standardized by microtiter plate ELISA with human secretory IgA. The product is tested for purity and specificity at final concentration by immunoelectrophoresis. The antibody is predominantly goat IgG; no trace of albumin is detected. It shows reactivity to human secretory IgA; cross-reactivity to other species may exist. Antibody activity to IgG, IgM, or non-immunoglobulin serum proteins is not present.

### EXPIRATION DATE:

5/31/2020

### DILUTION RECOMMENDATIONS:

1:500 - 1:4,000 for Tissue Staining  
1:1,000 - 1:8,000 for Immuno Blotting  
1:2,000 - 1:16,000 for ELISA/FIA

### RECONSTITUTION:

Reconstitute product with 2.0 ml of deionized or distilled water. Gentle swirling may be used to speed rehydration. Do not add sodium azide. Avoid vigorous shaking of the reconstituted material.

### STORAGE:

Store lyophilized product at 2-8°C. For long-term storage, aliquot and freeze reconstituted product at -20°C or lower.

### WARNING AND CAUTION:

This product is intended **FOR RESEARCH USE ONLY**, not for use in diagnostic or therapeutic procedures involving humans or animals.

### RELATED PRODUCT ORDERING INFORMATION:

Catalog #DescriptionSize  
55080 GOAT ANTI-HUMAN SECRETORY IGA2 ML  
55001 IGG GOAT ANTI-HUMAN SECRETORY IGA5 ML  
55046 F(AB')<sub>2</sub> GOAT ANTI-HUMAN SECRETORY IGA2 ML  
55905 PURIFIED HUMAN SECRETORY IGA5 MG  
55076 FITC GOAT ANTI-HUMAN SECRETORY IGA2 ML  
36315 DAB6x1ML  
36317 OPD6x1ML  
36318 TMB6x1ML  
36319 ABTS@6x1ML  
55984 NORMAL GOAT SERUM2 ML

**For fluorochrome conjugates, please see our catalog.**

ABTS is a registered trademark of Boehringer Mannheim GmbH.

**TO PLACE AN ORDER IN THE U.S., PLEASE CALL 800-854-0530  
OR TO OBTAIN TECHNICAL ASSISTANCE IN THE U.S., PLEASE CALL 800-279-5490  
OUTSIDE THE U.S., CONTACT YOUR LOCAL DISTRIBUTOR.**